

1. Details of Module and its structure

Module Detail	
Subject Name	Biology
Course Name	Biology 01 (Class XI, Semester - 1)
Module Name/Title	Kingdom Plantae - Bryophyta & Pteridophyta – Part 2
Module Id	kebo_10302
Pre-requisites	Basic Characteristics of Plants
Objectives	<p>After going through this lesson, the learners will be able to understand the following:</p> <ul style="list-style-type: none">• Adaptations in bryophytes that enable them live on land.• What are the three groups of bryophytes?• How do bryophytes reproduce?• How is vascular tissue important to pteridophytes ?• What are the characteristics of the four phyla of pteridophytes?• How pteridophytes reproduce?• Ecological role of bryophytes and pteridophytes.
Keywords	Bryophytes, Rhizoid, Gemma, Protonema, Antheridium, Archegonium, Vascular tissue, Sori, Rhizome, Pteridophytes, Hepaticopsida, Anthocerotopsida, Bryopsida, Lycophyta, Pterophyta, Sphenophyta

2. Development Team

Role	Name	Affiliation
National MOOC Coordinator (NMC)	Prof. Amarendra P. Behera	CIET, NCERT, New Delhi
Program Coordinator	Dr. Mohd. Mamur Ali	CIET, NCERT, New Delhi
Course Coordinator (CC) / PI	Dr. Sunita Farkya	DESM, NCERT, New Delhi
Course Co-Coordinator / Co-PI	Dr. Yash Paul Sharma	CIET, NCERT, New Delhi
Subject Matter Expert (SME)	Mr. Naqeeb Mehdi	Delhi Public School, Greater Noida
Review Team	Dr. K.V. Sridevi	RMSA Project Cell, NCERT, New Delhi

Table of Contents:

- 1) Introduction
- 2) Bryophyta-
 - a) Introduction to Bryophytes
 - b) Habitat and Distribution
 - c) Classification –
 - i) Mosses,
 - ii) Liverworts
 - iii) Hornworts
 - d) Life-cycle of bryophytes
 - e) Ecological Role
 - f) Economic Role
- 3) Pteridophyta –
 - a) Introduction to Pteridophyta
 - b) General Characteristics
 - c) Habitat and Distribution
 - d) Classification – Club Mosses & Horsetails
 - e) Economic importance of pteridophytes.
- 4) Summary

1) Introduction to Bryophytes and Pteridophytes –

The kingdom plantae includes eukaryotic, photosynthetic organisms. The kingdom plantae have traditionally been classified broadly into two sub-kingdoms cryptogamae and phanerogamae. As in the previous module we broadly studied the classification of plants with special relevance to the division of Thallophytes in this module we will learn about the other two divisions in cryptogamae. Bryophyta also known as “amphibians of the plant kingdom” and Pteridophyta which are often referred as the “first land plants or vascular cryptogams”.

2) **a) Bryophytes** – Bryophytes are simplest or the most primitive non-vascular plants, bryophytes have lifecycles that depend on water for fertilization in order to reproduce thus they are mostly found in moist places and thus they are also known as the ‘amphibians of the plant kingdom’. These plants completely lack vascular tissue, these plants can draw water by osmosis as they are only a few centimeters above the ground. Bryophytes include the various mosses, hornworts and liverworts that are found commonly growing in moist shaded areas in the hills. In this module we would learn about the general characteristics, habitat and distribution, classification, life-cycle, Ecological and economic importance of Bryophytes.

b) Habitat and distribution - Bryophytes are widely distributed throughout the world, from polar and alpine regions to the tropics. Bryophytes produce sperms that must swim through water in order to reach egg for fertilization thus bryophytes are not found in extremely arid region or saline water, although some are found in damp environments which are wet throughout the year within arid regions and some species are found on seashores above the intertidal zone. The tropical and subtropical latitudes show a great diversity of Bryophytes.

Bryophytes in India - Out of 16,236 species of 2523 species of bryophytes are found in India with about 629 endemic species and around 80 threatened species. (Source: Chapman, 2009; Singh & Dsah, 2014). Though the species of bryophytes are widely distributed within India the two hotspots of biodiversity one in the north east Himalaya and the other in the Western Ghats show richest diversity of Bryophytes within India.

c) Classification of Bryophytes- Bryophytes are generally classified into Mosses, Liverworts and Hornworts. Today we classify them into three phyla officially named i) Hepaticopsida

(Liverworts), ii) Anthocerotopsida (Hornworts) and iii) Bryopsida (Mosses)

i) **Hepaticopsida or Liverworts** – These are commonly called as liverworts as the gametophytic plant body may be thallose mostly foliose or leaf like with aseptate rhizoids gives an appearance of liver as seen in *Marchantia*, *Riccia*, *Pellia* etc. Sex organs are mostly found on the dorsal surface of the thallus. The male Antheridia are borne on antheridiophores which produce flagellated antherozoids or the sperms. The

female reproductive organs are archegonia are borne on specialized branches which are called as archegoniophores each archegonium is differentiated into neck and venter the egg is placed inside the venter. Post fertilization the egg becomes zygote (2N) which grows into a sporophyte also called sporangium . The body of the sporophyte is usually differentiated into bulbous foot, setae and an oval capsule as seen in *Marchantia*. Inside the capsule the sporogenous cells develop from the endothecium, the spore mother cell are diploid which divide through meiosis to give rise to haploid cells or the spores that later get dispersed by wind. Some of the liverworts also reproduce asexually by means of gemmae (Singular :gemma) are small multicellular reproductive structures. In some species like in *Marchantia*the gemma are produced in cuplike structures called gemma cup, the gemma can divide mitotically to give rise to new individual.

- ii) **Antocerotopsida or Hornworts** –like the liverworts, hornworts are generally found in soil that stays moist and damp throughout the year. Their gametophyte looks quite the same as liverworts i.e. thalloid with aseptate rhizoids. The sporophyte or sporangium is differentiated into bulbous foot, small meristematic seta and a long cylindrical capsule with pseudo elaters the sporophyte resemble small horn. Common example includes *Anthoceros* which is long and grows by means of basal meristematic zone. The sporogenous cells develop from amphitecium and endothecium forms sterile columella. The capsule dehisces irregularly basipetally by two valves and shows hygroscopic twisting. Some examples include *Anthoceros*, *Notothylus*, *Megaceros* etc.
- iii) **Bryopsida or Mosses** - These are the most common forms of bryophytes. They are commonly known as mosses. They grow in abundance in areas with water-bogs & swamps near streams and in rainforests. Mosses are able to tolerate low soil nutrient, low temperatures allowing them to grow in harsh environmental conditions. Thus mosses show a wide range of distribution with its presence even in the polar regions

the Arctic and the Antarctic. They usually form the green carpets or mats on moist soils, walls and tree trunks. The clumps of gametophyte grow together this predominant stage of the life cycle of a moss, the gametophyte consists of two stages. The first stage is the protonema stage, which develops directly from a spore. It is a creeping, green, branched and frequently filamentous stage. The second stage is the leafy stage, which develops from the secondary protonema as a lateral bud. They consist of upright, slender axes bearing spirally arranged leaves. They are attached to the soil through multicellular and branched obliquely septate rhizoids. This stage bears the sex organs. Vegetative reproduction in mosses is by fragmentation and budding in the secondary protonema. In sexual reproduction, the sex organs antheridia and archegonia are produced at the apex of the leafy shoots. After fertilisation, the zygote develops into a sporophyte, consisting of a foot, seta and capsule. The sporophyte in mosses is more elaborate than that in liverworts. The capsule contains spores. Spores are formed after meiosis. The mosses have an elaborate mechanism of spore dispersal. Common examples of mosses are *Funaria*, *Polytrichum* and *Sphagnum*.

d) Life-cycle of Bryophytes – The lifecycle of bryophytes consist of two distinct phases (i) Gametophytic phase and ii) Sporophytic phase. In Bryophytes the gametophyte is the dominant recognizable stage of the life-cycle that carries out most of the plants photosynthesis. The sporophyte which grows haploid spores grow at the top of the gametophyte plant. Let us first study the gametophytic plant body in bryophytes followed by the study of reproduction in bryophytes, sporophyte and the alternation of generation seen in bryophytes.

The Gametophytic plant body

The haploid gametophyte is dominant, long lived green and independent. The gametophyte are thalloid (i.e Not differentiated into true root, stems and true leaves) or leafy shoot having stem like central axis and leaf like appendages. The roots are completely absent and they are replaced by unicellular or multicellular thread like

rhizoids which help in absorbing moisture from the soil through osmosis. The vascular tissues i.e Xylem and Phloem are completely absent.

Reproduction- Bryophytes reproduce through both asexual and sexual methods of reproduction. Asexually bryophytes reproduce through fragmentation, adventitious branches, Gemmae etc.

The sexual reproduction in bryophytes is oogamous. i.e male gametes are smaller in comparison to large female gametes, also male sperms are flagellated and are called as antherozoids and female gametes are large and non-motile. The gametes are produced in multicellular, jacketed sex organs. The male reproductive are called antheridia and the female reproductive organs are called archegonia. Each antheridia is is differentiated stalk and body which produce fertile motile antherozoids. The archegonia is flask shaped differentiated into neck and venter, it is the venter that encloses the a large naked egg. During fertilization the male gamete released by the antheridia swim in water droplets which when comes in contact with archegonia fertilizes the egg. Since bryophytes require water for fertilization and fusion of gametes in order to reproduce thus they are also known as “The amphibians of the plant kingdom”.

The Sporophyte –The zygote that results in due to fertilization is a diploid cell that becomes the first cell of sporophytic generation that divides further to develop into a sporogonium. The sporogonium gets differentiated into foot seta and capsule, and the cells of the outer wall forms the calyptra that provides protective covering to developing sporogonium. The sporogonium asexually produces haploid spores as a result of meiosis that grows into gametophytic plant body.

Alternation of generation – The lifecycle of bryophytes show alternation of the two morphologically dissimilar phases in its lifecycle which is also known as alternation of generations. The alternation of generation in bryophytes is heteromorphic.

e) Ecological role of bryophytes –

- i) Formation of soil – Mosses and lichen plays a prominent role in the formation of soil as they grow on rocks and add its debris when it dies as organic matter which makes it suitable for higher forms plants to grow which in turns lead to breaking up of rock into piece which later weather leading to formation of soil.
- ii) Soil Conservation- Mosses play a very important role in binding the soil as it forms dense carpet like mats over the soil which it prevent soil erosion by water streams.
- iii) Pollution Indicators - they are sensitive indicators of atmospheric pollutants. Changes in the distributions of mosses (and lichens) are early-warning signal increase in atmospheric pollution.
- iv) Mineral Indicators – Certain forms of moss grows on soils which are rich in certain specific mineral. Eg Copper Moss growing over region is an indicator of soil being rich in copper.

f) Economic Importance of bryophytes –

- i) Use in nursery – *Sphagnum* moss are able to thrive in acidic water in bogs. Dried Sphagnum moss can absorb water many a times its own weight in water and thus acts sort of a natural sponge. Thus it is widely used by gardeners to keep cut plants moist during their propagation.
- ii) Peat - *Sphagnum* sometimes accumulates to form peat which is used by farmer to increase soils water holding capacity. and also is used in extraction of several chemical products like tar, paraffin, nitrates, tanning material etc.
- iii) Medicine – Certain forms of mosses help in extraction of antibiotic substances.

3) Introduction to Pteridophytes – After learning about the Bryophyta we would now be learning about seedless vascular plants called pteridophyte . These are cryptogams having sporophyte body and inconspicuous gametophyte. They are the first vascular plants and are often called as “botanical snakes ”. In the second of this module we would be learning about the general characteristics, habitat & distribution, classification, lifecycle, economic importance and ecological importance of pteridophytes.

a) **General Characteristics**—The term pteridophytes is derived from two greek word pteron which means feather and phyton which means plants. These plants usually recognized as ferns are diploid sporophyte which constitute the main and the dominant stage. The sporophyte is differentiated into roots, stems and leaves. The roots of pteridophyte are strong, creeping or underground stems called rhizome, the primary roots are short lived and is usually replaced by adventitious roots. The leaves may be small microphyllous as in case of *Lycopodium*, *Equisetum* or large macrophyllous as in *Pteridium*, *Pteris*. The vegetative parts of the plant contains vascular tissue that is xylem and phloem. In many plants the combination of the thick walls of xylem and lignin enables vascular plants grow and reach tall.

b) **Habitat and distribution** – They grow in variety of habitat but mostly terrestrial plants and thrive well in abundant moisture & shaded regions. However there are other forms of pteridophyte that are able to flourish in arid conditions. There are about 12000 species of pteridophyte of which 10.57% i.e. 1267 have been identified in India this includes 47 endemic and 414 threatened species. Most of the pteridophytes are found in Himalayas and the western ghats. The western ghats of Karnataka show high specie diversity amongst pteridophytes.

c) **Classification of pteridophytes** - Pteridophytes are classified into four groups –

i) **Psilophyta** – These are oldest land inhabiting plants. Rootless, rhizoids are present. These are homosporous most plants are fossils e.g. *Rhynia*, *Homeophyton* etc

ii) **Lycophyta**– commonly called club mosses or spike mosses. Roots, stem and leaves are present. *Lycopodium* and *Selaginella*.

***Lycopodium* Sp-** They have leaves which are small and microphyllous with a long creeping rhizome which give rise to aerial branches. These aerial branches bear sporangia in terminal cluster called strobili which are club shaped thus it also called as club moss.

Selaginella- They are also called as spike moss. Different species of *Selaginella* are widely distributed and are found in varied conditions from moist damp soil to temperate and xerophytic conditions. These plants have capacity to roll up to assume shape of ball during prolonged dry period and resume to usual form when dipped in water, such plants are also called '**resurrection plants**'. They also show heterospory which step is leading to evolution into '**seed habit**' i.e ability of plants to form seeds.

iii) Sphenophyta—At nodes whorls of small non photosynthetic leaves are present, Deposition of silica in stems make them rough in touch. Egs. *Equisetum*.

Equisetum – Commonly known as horsetails or scouring rushes. These are usually found growing in swampy soil growing along the banks of river etc. These plants usually may grow about a meter tall. They are called horsetail because its stems look similar to horse tails and due to presence of abrasive crystals of silica these are also known as scouring rushes, infact during colonial era horsetails were used to scour pots and pans.

iv) Pterophyta (FERNS) – They have fan like leaves called fronds, underground stem is in the form rhizome. They are found in shady and moist regions, are found very commonly in great numbers in tropical rainforests egs *Marselia*, *Pteris*, *Dryopteris*.

d) Life-Cycle of Ferns – The dominant phase in ferns as discussed earlier is the sporophyte which is actually diploid. These plants develop spores on the underside of their fronds in structures which are called as sporangia. The clusters of sporangia are called as sori. The spores are meant for reproduction and germinate to give rise to haploid gametophyte, called prothallus. The homosporous pteridophyte produce bisexual (monoecious) gametophyte whereas hetrosporous pteridophytes produce unisexual (dioecious) gametophytes. When the spores germinate they developed into a small gametophyte that produces root like rhizoids, the gametophyte though is small but still it grows independent of the sporophyte. The gametophyte produces archegonia and antheridia on the underside of the gametophyte. The fertilization is oogamous and requires water as it was in bryophytes to develop into a diploid zygote that develops into sporophyte.

e) Economic role of pteridophytes –

- i. **Soil Binding**–Pteridophytes binds the soil all along hill slopes thus preventing soil erosion.
- ii. **Food**- Pteridophytes constitutes a good source of food for animals.Sporocarps of Marselia are cooked and eaten by certain tribes.
- iii. **Scouring**- Stems of equisetum helps in scouring of utensils and polishing of metals.
Egs. *Equisetum*.
- iv. **Nitrogen fixing** – Some ferns eg Water fern Azolla form symbiotic association with nitrogen fixing cyanobacteriumare used as bio fertilisers in paddy fields.
- v. **Medicines** – Many antihelminthic drug is obtained from the rhizome and petioles of the fern- *Dryopteris*. *Lycopodium* is used in the treatment of rheumatism and disorder of lungs and kidneys. *Adiantum* roots are used to cure throat infections.
- vi. **Ornamental** – Ferns are known for their attractive graceful foliage.

4) Summary - In this module we learnt about two subdivisions under cryptogams i.e Bryophytes and Pteridophytes. Lets recall what we had learn about each of them :

BRYOPHYTES - Bryophytes (division Bryophyta) are the simplest of the terrestrial plants that lack complex tissue organization, yet they show considerable diversity in form and ecology. The division Bryophytaare classified into three phyla-

1. The mosses (class Bryopsida)
2. The liverworts (class Hepatopsida)
3. The hornworts (class Anthocerotopsida)

The bryophytes show an alternation of generations between the independent gametophyte generation, which produces the sperm and eggs, and the dependent sporophyte generation, which produces the spores. These are known as amphibians of plant kingdom as they need presence of water for fertilization of gametes.

Bryophytes form mats, spongy carpets. Bryophytes especially Mosses which play an important key role in forming ecological communities in environment as they help in formation of soil
Bryophytes are very sensitive to pollution and are bio-indicators of environment.

Ferns or Pteridophytes are primitive vascular plants that don't have seeds. In pteridophytes the main plant body is a diploid sporophyte which is differentiated into stem, roots and leaves. Pteridophytes are classified into two groups (i) Psilophyta (ii) Lycophyta (Club moss and Spike moss) (iii) Sphenophyta (Horsetails) (iv) Pterophyta (Fern). The sporophyte is the dominant phase and the spores are meant for reproduction which germinate to give rise to haploid gametophyte, called prothallus. The gametophyte produces archegonia and antheridia on the underside of the gametophyte. The fertilization is oogamous and requires water as it was in bryophytes to develop into a diploid zygote that develops into sporophyte.

Pteridophytes have important economic importance as they are used as food, fixing nitrogen, medicine, ornamental purpose and also helps in binding the soil.