

## 1. Details of Module and its structure

Module Detail	
Subject Name	Chemistry
Course Name	Chemistry 01 (Class XI, Semester 01)
Module Name/Title	Chemical bonding and molecular structure: Part1
Module Id	kech_10401
Pre-requisites	Knowledge about atoms, molecules, atomic structure and electronic configuration
Objectives	After going through this module, the learners will be able to: <ul style="list-style-type: none"><li>• Understand kössel-Lewis approach to chemical bonding</li><li>• Explain the octet rule and its limitations</li><li>• Draw Lewis structures of simple molecules</li><li>• Explain the formation of different types of bonds</li><li>• Concept of formal charge</li></ul>
Keywords	Lewis Structure, Octet Rule, Covalent and Electrovalent Bond, Formal Charge

## 2. Development Team

Role	Name	Affiliation
National MOOC Coordinator (NMC)	Prof. Amarendra P. Behera	CIET, NCERT, New Delhi
Program Coordinator	Dr. Mohd. Mamur Ali	CIET, NCERT, New Delhi
Course Coordinator (CC) / PI	Prof. R. K. Parashar	DESM, NCERT, New Delhi
Course Co-Coordinator / Co-PI	Dr. Aerum Khan	CIET, NCERT, New Delhi
Subject Matter Expert (SME)	Preeti Kiran	Kendriya Vidyalaya No1, AFS, Hindan, GZB
Review Team	Dr. Sulekh Chandra	Zakir Hussain College, New Delhi
	Dr. Aerum Khan	CIET, NCERT, New Delhi

---

## Table of Contents:

1. Introduction: Chemical bonding and molecular structure
2. Kössel-Lewis approach to chemical bonding
3. Octet rule
4. Covalent and Electrovalent bond
5. Formal charge
6. Limitations of the Octet Rule
7. Summary

### 1. Introduction: Chemical bonding and molecular structure

Matter is made up of one or different type of elements. Under normal conditions no other element exists as an independent atom in nature, except noble gases. Most of these atoms exist as molecules which are cluster or aggregates of atoms or ions of same or different elements having a distinct existence and its own characteristic properties. Obviously there must be some force which holds these constituent atoms together in the molecules.

**The attractive force which holds various constituents (atoms, ions, etc.) together in different chemical species is called a chemical bond.**

Since the formation of chemical compounds takes place as a result of combination of atoms of various elements in different ways, it raises many questions.

- Why do atoms combine?
- Why are only certain combinations possible?
- Why do some atoms combine while certain others do not?
- Why do atoms have fixed combining capacity?
- Why do molecules possess definite shapes?

To answer such questions different theories and concepts have been put forward from time to time. These are

- Kössel-Lewis approach,
- Valence Shell Electron Pair Repulsion (VSEPR) Theory,
- Valence Bond (VB) Theory and
- Molecular Orbital (MO) Theory.

The evolution of various theories of valence and the interpretation of the nature of chemical bonds have closely been related to the developments in the understanding of the structure of atom, the electronic configuration of elements and the periodic table.

---

Every system tends to be more stable and bonding is nature's way of lowering the energy of the system to attain stability.

## 2. Kössel-Lewis approach to chemical bonding

To explain the formation of chemical bond in terms of electrons, a number of attempts were made, but it was only in 1916 when Kössel and Lewis succeeded independently provide logical explanation of valence which was based on the inertness of noble gases.

The atoms of noble gases except helium have eight electrons in their valence shell. The valence shell electronic configuration of noble gas atoms is (except Helium) is  $ns^2np^6$  which represents the stable configuration and corresponds to maximum stability. Due to stable configuration, the noble gases neither have any tendency to gain or lose electrons and therefore their combining capacity or valency is zero. Their inertness is so prominent that they exist as monoatomic gaseous atoms.

All atoms other than noble gases have less than eight electrons in their outermost shell i.e they don't have stable electronic configuration so they combine with each other or other atoms to achieve stable noble gas electronic configuration. ( $ns^2np^6$  or  $1s^2$ ) which is the state of minimum energy and maximum stability. Atoms combine with each other only if the process leads to lowering of energy.


Lewis depicted the atom in terms of a positively charged 'Kernel' (the nucleus plus the inner electrons) and the outer shell that could accommodate a maximum of eight electrons. He, further assumed that these eight electrons occupy the corners of a cube which surround the 'Kernel'. Thus the single valence shell electron of sodium would occupy one corner of the cube, while in the case of a noble gas all the eight corners would be occupied. This octet of electrons, represents a particularly stable electronic arrangement.

Lewis postulated that atoms achieve the stable octet when they are linked by chemical bonds. In the case of sodium and chlorine, this can happen by the transfer of an electron from sodium to chlorine thereby giving the  $Na^+$  and  $Cl^-$  ions.

In the case of other molecules like  $Cl_2$ ,  $H_2$ ,  $F_2$ , etc., the bond is formed by the sharing of a pair of electrons between the atoms. In the process each atom attains a stable outer octet of electrons.

**Lewis Symbols:** In the formation of a molecule, only the outer shell electrons take part in chemical combination and they are known as valence electrons. The inner shell electrons are well protected and are generally do not participate in the combination process.


G.N. Lewis, an American chemist introduced simple notations to represent valence electrons in an atom. These notations are called Lewis symbols. For example, the Lewis symbols for the elements of second period are as under :





**Significance of Lewis Symbols:** The number of dots around the symbol represents the number of valence electrons. This number of valence electrons helps to calculate the common or group valence of the element. The group valence of the elements is generally either equal to the number of dots in Lewis symbols or 8 minus the number of dots or valence electrons.

Kössel, in relation to chemical bonding, drew attention to the following facts:

- In the periodic table, the highly electronegative halogens and the highly electropositive alkali metals are separated by the noble gases;
- The formation of a negative ion from a halogen atom and a positive ion from an alkali metal atom is associated with the gain and loss of an electron by the respective atoms;
- The negative and positive ions thus formed attain stable noble gas electronic configurations. The noble gases (with the exception of helium which has a duplet of electrons) have a particularly stable outer shell configuration of eight (octet) electrons,  $ns^2np^6$ .
- The negative and positive ions are stabilized by electrostatic attraction.
- For example, the formation of NaCl from sodium and chlorine, according to the above scheme, can be explained as:



Similarly the formation of  $\text{CaF}_2$  may be shown as:





The bond formed, as a result of the electrostatic attraction between the positive and negative ions was termed as the electrovalent bond. The electrovalence is thus equal to the number of unit charge(s) on the ion. Thus, calcium is assigned a positive electrovalence of two, while chlorine a negative electrovalence of one.

Kössel's postulations provide the basis for the modern concepts regarding ion-formation by electron transfer and the formation of ionic crystalline compounds. His views have proved to be of great value in the understanding and systematisation of the ionic compounds. At the same time he did recognise the fact that a large number of compounds did not fit into these concepts.

### 3. Octet Rule

Kössel and Lewis in 1916 developed an important theory of chemical combination between atoms known as electronic theory of chemical bonding. According to this, "Atoms can combine either by transfer of valence electrons from one atom to another (gaining or losing) or by sharing of valence electrons in order to have an octet in their valence shells. This is known as octet rule".


### 4. Covalent and Electrovalent Bond

The Lewis theory was the first to explain formation of a covalent bond in terms of electrons that was generally accepted. If two electrons are shared between two atoms, this constitutes a bond and binds the atoms together. For many lighter atoms a stable arrangement is attained when the atom is surrounded by eight electrons. This octet can be made up from some electrons which are 'totally owned' and some electrons which are 'shared'. Thus atoms continue to form bonds (single and multiple) until they complete an octet of electrons. This is called the 'octet rule'. The octet rule explains the observed valencies in a large number of cases.

**Langmuir (1919) refined the Lewis postulations** by abandoning the idea of the stationary cubical arrangement of the octet, and by **introducing the term covalent bond**.

The Lewis-Langmuir theory can be understood by considering the formation of the chlorine molecule,  $\text{Cl}_2$ .

The Cl atom with electronic configuration,  $[\text{Ne}]3s^2 3p^5$ , is one electron short of the argon configuration. The formation of the  $\text{Cl}_2$  molecule can be understood in terms of the sharing of a pair of electrons between the two chlorine atoms, each chlorine atom contributing one electron to the shared pair. In the process both chlorine atoms attain the outer shell octet of the nearest noble gas (i.e., argon).




Cl – Cl      *Covalent bond between two Cl atoms*

**The dots represent electrons. Such structures are referred to as Lewis dot structures.**

The Lewis dot structures can be written for other molecules also, in which the combining atoms may be identical or different. The important conditions being that:

- Each bond is formed as a result of sharing of an electron pair between the atoms.
- Each combining atom contributes at least one electron to the shared pair.
- The combining atoms attain the outershell noble gas configurations as a result of the sharing of electrons.
- Thus in water and carbon tetrachloride molecules, formation of covalent bonds can be represented as:




H atoms attain a duplet of electrons and O, the octet


Each of the four Cl atoms along with the C atom attains octet of electrons

Thus, when two atoms share one electron pair they are said to be joined by a single covalent bond. In many compounds we have multiple bonds between atoms. The formation of multiple bonds envisages sharing of more than one electron pair between two atoms.



If two atoms share two pairs of electrons, the covalent bond between them is called a double bond. For example, in the carbon dioxide molecule, we have two double bonds between the carbon and oxygen atoms.



Similarly in ethene molecule the two carbon atoms are joined by a double bond.



When combining atoms share three electron pairs as in the case of two nitrogen atoms in the N<sub>2</sub> molecule and the two carbon atoms in the ethyne molecule, a triple bond is formed.



### Lewis Representation of Simple Molecules (the Lewis Structures)

---

The Lewis dot structures provide a picture of bonding in molecules and ions in terms of the shared pairs of electrons and the octet rule. While such a picture may not explain the bonding and behaviour of a molecule completely, it does help in understanding the formation and properties of a molecule to a large extent.









**Writing of Lewis dot structures of molecules** is, therefore, very useful. The Lewis dot structures can be written by adopting the following steps:

- The total number of electrons required for writing the structures are obtained by adding the valence electrons of the combining atoms.
- For example, in the CH<sub>4</sub> molecule there are eight valence electrons available for bonding (4 from carbon and 4 from the four hydrogen atoms).
- For anions, each negative charge would mean addition of one electron.
- For cations, each positive charge would result in subtraction of one electron from the total number of valence electrons.
- For example, for the CO<sub>3</sub><sup>2-</sup> ion, the two negative charges indicate that there are two additional electrons than those provided by the neutral atoms. For NH<sub>4</sub><sup>+</sup> ion, one positive charge indicates the loss of one electron from the group of neutral atoms.
- Knowing the chemical symbols of the combining atoms and having knowledge of the skeletal structure of the compound (known or guessed intelligently), it is easy to distribute the total number of electrons as bonding shared pairs between the atoms in proportion to the total bonds.
- In general the least electronegative atom occupies the central position in the molecule/ion. For example in the NF<sub>3</sub> and CO<sub>3</sub><sup>2-</sup>, nitrogen and carbon are the central atoms whereas fluorine and oxygen occupy the terminal positions.
- After accounting for the shared pairs of electrons for single bonds, the remaining electron pairs are either utilized for multiple bonding or remain as the lone pairs. The basic requirement being that each bonded atom gets an octet of electrons.

Lewis representations of a few molecules/ ions are given in Table 1.



Table 1: The Lewis Representation of some molecules.

Molecule/Ion	Lewis Representation	Lewis Representation
H <sub>2</sub>	H : H*	H – H
O <sub>2</sub>	:Ö::Ö:	:Ö=Ö:
O <sub>3</sub>		
NF <sub>3</sub>		
CO <sub>3</sub> <sup>2-</sup>		
HNO <sub>3</sub>		

### Problem 1

Write the Lewis dot structure of CO molecule.


#### Solution

**Step 1.** Count the total number of valence electrons of carbon and oxygen atoms. The outer (valence) shell configurations of carbon and oxygen atoms are:  $2s^2 2p^2$  and  $2s^2 2p^4$ , respectively.


The valence electrons available are  $4 + 6 = 10$ .

**Step 2.** The skeletal structure of CO is written as: C O

**Step 3.** Draw a single bond (one shared electron pair) between C and O and complete the octet on O, the remaining two electrons are the lone pair on C.



This does not complete the octet on carbon and hence we have to resort to multiple bonding (in this case a triple bond) between C and O atoms. This satisfies the octet rule condition for both atoms.



### Problem 2

Write the Lewis structure of the nitrite ion, NO<sub>2</sub><sup>-</sup>.

## Solution


**Step 1.** Count the total number of valence electrons of the nitrogen atom, the oxygen atoms and the additional one negative charge (equal to one electron).




$$5 + (2 \times 6) + 1 = 18 \text{ electrons}$$

**Step 2.** The skeletal structure of  $\text{NO}_2^-$  is written as : O N O

**Step 3.** Draw a single bond (one shared electron pair) between the nitrogen and each of the oxygen atoms completing the octets on oxygen atoms. This, however, does not complete the octet on nitrogen if the remaining two electrons constitute lone pair on it.



Hence we have to resort to multiple bonding between nitrogen and one of the oxygen atoms (in this case a double bond). This leads to the following Lewis dot structures.



## 5. Formal Charge

Lewis dot structures, in general, do not represent the actual shapes of the molecules. In case of polyatomic ions, the net charge is possessed by the ion as a whole and not by a particular atom. It is, however, feasible to assign a formal charge on each atom.

“The formal charge of an atom in a polyatomic molecule or ion may be defined as the difference between the number of valence electrons of that atom in an isolated or free state and the number of electrons assigned to that atom in the Lewis structure.”


- A **formal charge (FC)** is the charge assigned to an atom in a molecule, assuming that electrons in all chemical bonds are shared equally between atoms, regardless of relative electronegativity. When determining the best Lewis structure (or predominant resonance structure) for a molecule, the structure is chosen such that the formal charge on each of the atoms is as close to zero as possible.
- The formal charge of any atom in a molecule can be calculated by the following equation:

$$FC = V - N - B/2$$

where **V** is the number of valence electrons of the neutral atom in isolation (in its ground state); **N** is the number of non-bonding valence electrons on this atom in the molecule; and **B** is the total number of electrons shared in bonds with other atoms in the molecule.

The counting is based on the assumption that the atom in the molecule owns one electron of each shared pair and both the electrons of a lone pair.


Let us consider the ozone molecule ( $O_3$ ). The Lewis structure of  $O_3$  may be drawn as:



The atoms have been numbered as 1, 2 and 3. The formal charge on:

- The central O atom marked 1  
 $= 6 - 2 - 1/2(6) = +1$
- The end O atom marked 2  
 $= 6 - 4 - 1/2(4) = 0$
- The end O atom marked 3  
 $= 6 - 6 - 1/2(2) = -1$

Hence, we represent  $O_3$  along with the formal charges as follows:



---

We must understand that formal charges do not indicate real charge separation within the molecule. Indicating the charges on the atoms in the Lewis structure only helps in keeping track of the valence electrons in the molecule.

Formal charges help in the selection of the lowest energy structure from a number of possible Lewis structures for a given species.

Generally the lowest energy structure is the one with the smallest formal charges on the atoms. The formal charge is a factor based on a pure covalent view of bonding in which electron pairs are shared equally by neighbouring atoms.

## 6. Limitations of the Octet Rule

The octet rule, though useful, is not universal. It is quite useful for understanding the structures of most of the organic compounds and it applies mainly to the second period elements of the periodic table. There are three types of exceptions to the octet rule.

### The incomplete octet of the central atom


In some compounds, the number of electrons surrounding the central atom is less than eight. This is especially the case with elements having less than four valence electrons. Examples are LiCl, BeH<sub>2</sub> and BCl<sub>3</sub>.



Li, Be and B have 1,2 and 3 valence electrons only. Some other such compounds are AlCl<sub>3</sub> and BF<sub>3</sub>.

### Odd-electron molecules

In molecules with an odd number of electrons like nitric oxide, NO and nitrogen dioxide, NO<sub>2</sub>, the octet rule is not satisfied for all the atoms




### The expanded octet

The octet rule is also broken where atoms have an extra energy level which is close in energy to the p level .


For example Elements in and beyond the third period of the periodic table have, apart from 3s and 3p orbitals, 3d orbitals also available for bonding. In a number of compounds of these elements there are more than eight valence electrons around the central atom. This is termed as the expanded octet. Obviously the octet rule does not apply in such cases.

Some of the examples of such compounds are:  $\text{PF}_3$  obeys the octet rule, but  $\text{PF}_5$  does not.  $\text{PF}_5$  has ten outer electrons, and uses one  $3s$ , three  $3p$  and one  $3d$  orbitals.



Any compound with more than four covalent bonds must break the octet rule, and these violations become increasingly common in elements after the first two periods of eight elements in the periodic table.

Interestingly, sulphur also forms many compounds in which the octet rule is obeyed. In sulphur dichloride, the S atom has an octet of electrons around it.



### Other drawbacks of the octet theory

- It is clear that octet rule is based upon the chemical inertness of noble gases. However, some noble gases (for example xenon and krypton) also combine with oxygen and fluorine to form a number of compounds like  $\text{XeF}_2$ ,  $\text{KrF}_2$ ,  $\text{XeOF}_2$  etc.,
- This theory does not account for the shape of molecules.
- It does not explain the relative stability of the molecules being totally silent about the energy of a molecule.

# Lewis Dot Structures

- These symbols are called Lewis symbols →
- We generally place the electrons on four sides of a square around the element symbol.

Element	Electron Configuration	Electron-Dot Symbol
Li	[He]2s <sup>1</sup>	Li·
Be	[He]2s <sup>2</sup>	·Be·
B	[He]2s <sup>2</sup> 2p <sup>1</sup>	·B·
C	[He]2s <sup>2</sup> 2p <sup>2</sup>	·C·
N	[He]2s <sup>2</sup> 2p <sup>3</sup>	·N·
O	[He]2s <sup>2</sup> 2p <sup>4</sup>	·O·
F	[He]2s <sup>2</sup> 2p <sup>5</sup>	·F·
Ne	[He]2s <sup>2</sup> 2p <sup>6</sup>	·Ne·

## 4 step process: drawing lewis dot structure

**Step 1:** arrange atoms into a skeleton molecule–

- Central atom of lower electronegativity.
- The element written first in the formula is usually the central atom

**Step 2:** add up the valence electrons of all atoms

**Step 3:** draw single bonds and Subtract 2 electrons for each single bond from total counted in step 2


**Step 4:** deal out remaining electrons – create double or triple bond -satisfy octet rule (except for hydrogen)

Check: total electrons used in the Lewis structure should be equal to total valence electrons in step 1

Remember while drawing lewis structure:

- Hydrogen and halogens form single bonds
- Oxygen family need atleast two bonds to complete octet so they form single or double bond.
- Nitrogen family need three bonds so they form single , double or triple bond.
- Carbon family needs four bonds.

### The steps in converting a molecular formula into a Lewis structure.



## 7. Summary

The Lewis structure was named after Gilbert N. Lewis, who introduced it in his 1916 article. Lewis dot structure represent the molecular geometry and valence shell electrons of a molecule. Also known as Lewis diagrams/formulas/symbols. Most useful for atoms and molecules of period 2 and 3. It includes only valence electrons. Lewis structures extend the concept of the electron dot diagramly adding lines between atoms to represent shared pairs in a chemical bond. Lewis structures show each atom and its position in the structure of the molecule using its chemical symbol. Lines are drawn between atoms that are bonded to one another (pairs of dots can be used instead of lines). Excess electrons that form lone pairs are represented as pairs of dots, and are placed next to the atom.