1. Details of Module and Its Structure

Module Detail			
Subject Name	Chemistry		
Course Name	Chemistry 03 (Class XII, Semester 01)		
Module Name/Title	Chemical Kinetics: Part 2		
Module Id	lech_10402		
Pre-requisites	Chemical Reaction, concentration terms, chemical kinetics, rate of reaction, instantaneous rate of reaction, units of rate of reaction.		
Objectives	 After studying this module, the student would be able to: Discuss the dependence of rate of reaction on concentration Define rate law and rate constant Differentiate between the molecularity and the order of a reaction 		
Keywords	Rate of the reaction, factors affecting the rate of the reaction, effect of concentration on the rate of the reaction, rate law, rate constant, order of the reaction, molecularity of the reaction		

2. Development Team

Role	Name	Affiliation	
National MOOC Coordinator (NMC)	Prof. Amarendra P. Behera	CIET, NCERT, New Delhi	
Program Coordinator	Dr. Mohd. Mamur Ali	CIET, NCERT, New Delhi	
Course Coordinator (CC) / PI	Prof. Alka Mehrotra Prof. R. K. Parashar	DESM, NCERT, New Delhi DESM, NCERT, New Delhi	
Course Co-Coordinator / Co-PI	Dr. Aerum Khan	CIET, NCERT, New Delhi	
Subject Matter Expert (SME)	Dr. Komal S. Khatri	G. B. Pant Institute of Polytechnic, Okhla Phase II, New Delhi	
Review Team	Dr. Amirtha Anand	Dept. of Chemistry, Maitreyi College, New Delhi- 110021	

Table of Contents :

- 1. Factors affecting the rate of the reaction
- 2. Effect of concentration on the rate of the reaction
- 3. Rate law and rate constant
- 4. Order of a reaction
- 5. Molecularity of a reaction
- 6. Summary

1. Factors affecting the rate of the reaction: Rate of a reaction is influenced by a number of factors as follows:

- Experimental conditions such as concentration of reactants (pressure in case of gases);
- Temperature of the reactants or products;
- Nature of the reacting substances;
- Presence of catalyst.

2. Effect of concentration on the rate of the reaction: The rate of a chemical reaction at a given temperature may depend on the concentration of one or more reactants and products. It is found that the concentrations of the reactants decreases while the concentrations of the products increases with passage of time (Fig. 1).



Fig 1. Variation of concentration with passage of time

(Source:<u>https://upload.wikimedia.org/wikipedia/commons/thumb/2/24/ChemicalEquilibrium.</u> svg/300px-ChemicalEquilibrium.svg.png)

The representation of the rate of a reaction in terms of concentration of the reactants is known as rate law. It is also known as rate equation or rate expression.

3. Rate law and rate constant: In the first module, we have calculated the average rate of the reaction for hydrolysis of butyl chloride with passage of time. We found that the rate of a reaction decreases with the passage of time as the concentration of reactants decrease.

Conversely, rates generally increase when reactant concentrations increase. So, rate of a reaction depends upon the concentration of reactants.

Consider a general reaction

 $aA + bB \rightarrow cC + dD$

where a, b, c and d are the stoichiometric coefficients of reactants and products.

The rate expression for this reaction is

$$Rate \propto \left[A\right]^{x} \left[B\right]^{y} \tag{1}$$

where exponents x and y may or may not be equal to the stoichiometric coefficients (a and b) of the reactants. Above equation can also be written as

$$Rate = k[A]^{x}[B]^{y}$$
⁽²⁾

$$\frac{-d[R]}{dt} = k[A]^{x}[B]^{y}$$
(3)

This form of equation (3) is known as differential rate equation where k is a proportionality constant called rate constant. The equation like (1), which relates the rate of a reaction to concentration of reactants, is called rate law or rate expression. Thus, rate law is the expression in which reaction rate is given in terms of molar concentration of reactants with each term raised to some power, which may or may not be same as the stoichiometric coefficient of the reacting species in a balanced chemical equation.

Rate constant (k) also known as velocity constant or specific reaction rate, is defined as the rate of the reaction when concentration of each reactant is taken as unity.

As $[A] = [B] = 1 \mod L^{-1}$, *Rate* = k

Thus, the rate constant gives an idea about the speed of the reaction, i.e. a reaction with higher value of rate constant proceeds at faster rate. The rate constant depends upon temperature while independent of concentration of the reacting species.

The rate law expression can be explained using the following example:

$$2 \operatorname{NO}(g) + \operatorname{O}_2(g) \rightarrow 2 \operatorname{NO}_2(g)$$

We can measure the rate of this reaction as a function of initial concentrations either by keeping the concentration of one of the reactants constant and changing the concentration of the other reactant or by changing the concentration of both the reactants. The following results are obtained (Table 1).

Experiment	Initial [NO] / mol L ⁻	Initial $[O_2]$ / mol L^{-1}	Initial rate of formation of NO ₂ / mol L ⁻¹ s ⁻¹
1	0.30	0.30	0.096
2	0.60	0.30	0.384
3	0.30	0.60	0.192
4	0.60	0.60	0.768

Table 1. Initial rate of formation of NO_2

It is obvious, after looking at the results, that when the concentration of NO is doubled and that of O_2 is kept constant then the initial rate increases by a factor of four from 0.096 to 0.384 mol L⁻¹s⁻¹. This indicates that the rate depends upon the square of the concentration of NO (i.e. rate \propto [NO]²). When concentration of NO is kept constant and concentration of O_2 is doubled the rate also gets doubled indicating that rate depends on concentration of O_2 to the first power (i.e. rate \propto [O₂]). Hence, the rate equation for this reaction will be Rate = k [NO]²[O₂]

The differential form of this rate expression is given as

$$\frac{-d[R]}{dt} = k [NO]^2 [O_2]$$

Now, we observe that for this reaction, in the rate equation derived from the experimental data, the exponents of the concentration terms are the same as their stoichiometric coefficients in the balanced chemical equation.

Some other examples are given below:

- 1. CHCl₃ + Cl₂ \rightarrow CCl ₄ + HCl; the experimental rate expression is rate = k [CHCl₃] [Cl₂]^{1/2}
- 2. $CH_3COOC_2H_5 + H_2O \rightarrow CH_3COOH + C_2H_5OH$; the experimental rate expression is rate = k [CH_3COOC_2H_5]^1 [H_2O]^0

In these reactions, the exponents of the concentration terms are not the same as their stoichiometric coefficients. Thus, we can say that:

Rate law for any reaction cannot be predicted by merely looking at the balanced chemical equation, i.e., theoretically but must be determined experimentally.

Before going on further, it is very important to know the difference between the rate of the reaction and the rate constant of the reaction. As discussed above, the rate of the reaction is expressed as the speed of the reaction at which reactants are converted into products, while the rate constant is the proportionality constant in the rate law expression. Also, the rate of the reaction depends upon the concentration of the reactant species at a particular moment of time, while rate constant is equal to the rate of the reaction when concentration terms are unity at that moment of time. The rate of the reaction generally decreases with progress of the reaction; on the other hand, the rate constant is a constant value for a reaction at given conditions.

4. Order of a Reaction: In the rate equation (2)

Rate = $k [A]^x [B]^y$

where x and y indicate how sensitive the rate is to the change in concentration of A and B. Sum of these exponents, i.e., x + y in (2) gives the overall order of a reaction whereas x and y represent the order with respect to the reactants A and B respectively.

Hence, the sum of powers of the concentration of the reactants in the rate law expression is called the order of that chemical reaction.

Order of a reaction can be 0, 1, 2, 3 and even a fraction. A zero order reaction means that the rate of reaction is independent of the concentration of reactants.

A balanced chemical equation never gives us a true picture of how a reaction takes place since rarely a reaction gets completed in one step. The reactions taking place in one step are called elementary reactions. When a sequence of elementary reactions (called mechanism) gives us the products, the reactions are called complex reactions. These may be consecutive reactions (e.g., oxidation of ethane to CO₂ and H₂O passes through a series of intermediate steps in which alcohol, aldehyde and acid are formed), reverse reactions and side reactions (e.g., nitration of phenol yields o-nitrophenol and p-nitrophenol).

Units of rate constant: For a general reaction,

 $aA + bB \rightarrow cC + dD$

Rate = $k [A]^{x} [B]^{y}$

where x + y = n = order of the reaction
$$k = \frac{Rate}{[A]^{x}[B]^{y}}$$

= $\frac{concentration}{time} \times \frac{1}{(concentration)^{n}}$
= $(concentration)^{1-n} \times time^{-1}$

(where [A] = [B])

Taking SI units of concentration, mol L^{-1} and time, sec, the units of k for different reaction order are listed in Table 2.

Reaction	Order	Units of rate constant
Zero order reaction	0	$\frac{mol L^{-1}}{s} \times \frac{1}{(mol L^{-1})^{0}} = mol L^{-1} s^{-1}$
First order reaction	1	$\frac{mol L^{-1}}{s} \times \frac{1}{(mol L^{-1})^{1}} = s^{-1}$
Second order reaction	2	$\frac{mol L^{-1}}{s} \times \frac{1}{(mol L^{-1})^2} = mol^{-1} L s^{-1}$
n th order reaction	n	$\frac{mol L^{-1}}{s} \times \frac{1}{(mol L^{-1})^n} = (mol L^{-1})^{1-n} s^{-1}$

Table 2. Units of rate constant

Example 1: Calculate the overall order of a reaction which has the rate expression:

(a) Rate = k $[A]^{1/2} [B]^{3/2}$

(b) Rate = $k [A]^{3/2} [B]^{-1}$

Solution: (a) Rate = $k [A]^x [B]^y$

Order of the reaction = x + y

Here, $x = \frac{1}{2}$ and $y = \frac{3}{2}$

So order = 1/2 + 3/2 = 2, i.e., the reaction is a second order reaction.

(b) Here, x = 3/2 and y = -1

So order = 3/2 + (-1) = 1/2, i.e., the reaction is a half order reaction order.

Example 2: Identify the reaction order from each of the following rate constants.

- (i) $k = 2.3 \times 10^{-5} \text{ L mol}^{-1} \text{ s}^{-1}$
- (ii) $k = 3 \times 10^{-4} s^{-1}$

Solution: (i) From the table 2, we know that the unit of second order rate constant is $mol^{-1} L$ s⁻¹. In this case k = $2.3 \times 10^{-5} L mol^{-1} s^{-1}$ and hence it represents a second order reaction.

(ii) Again from the table 2, the unit of a first order rate constant is s^{-1} . Here, $k = 3 \times 10^{-4} s^{-1}$ and thus it represents a first order reaction.

Example 3: Find out the order of the reaction and units of rate constant in the following reactions. Also state the order with respect to each of the reactants.

(a) $CH_3CHO \rightarrow CH_4 + CO$; rate = k $[CH_3CHO]^{3/2}$ (b) CO + Cl₂ \rightarrow COCl₂; rate = k [CO]² [Cl₂]^{1/2} (c) NO + O₂ \rightarrow 2 NO₂; rate = k [NO]² [O₂] **Solution:** (a) Rate = $k [CH_3CHO]^{3/2}$ Order with respect to $CH_3CHO = 3/2 = 1.5$ Overall order of the reaction = 3/2 = 1.5Units of rate constant, $k = (mol L^{-1})^{1-n} s^{-1}$ Here, n = 1.5, therefore, units of k = (mol L^{-1})^{1-1.5} s⁻¹ = mol^{-1/2} $L^{1/2}$ s⁻¹ (b) Rate = k $[CO]^2 [Cl_2]^{1/2}$ Order with respect to CO = 2 and order with respect to $Cl_2 = \frac{1}{2} = 0.5$ Overall order of the reaction = $2 + \frac{1}{2} = \frac{21}{2} = \frac{5}{2} = 2.5$ Units of rate constant, $k = (mol L^{-1})^{1-n} s^{-1}$ Here, n = 2.5, therefore, units of k = (mol L^{-1})^{1-2.5} s⁻¹ = mol^{-3/2} $L^{3/2}$ s⁻¹ (c) Rate = $k [NO]^2 [O_2]$ Order with respect to NO = 2 and order with respect to $O_2 = 1$ Overall order of the reaction = 2 + 1 = 3Units of rate constant, $k = (mol L^{-1})^{1-n} s^{-1}$ Here, n = 3, therefore, units of k = $(mol L^{-1})^{1-3} s^{-1} = mol^{-2} L^2 s^{-1}$

5. Molecularity of a Reaction: Another property of a reaction called molecularity helps in understanding its mechanism. The number of reacting species (atoms, ions or molecules) taking part in an elementary reaction, which must collide simultaneously in order to bring about a chemical reaction is called molecularity of a reaction. The reaction can be unimolecular when one reacting species is involved, for example, decomposition of ammonium nitrite.

 $NH_4NO_2 \rightarrow N_2 + 2H_2O$

Bimolecular reactions involve simultaneous collision between two species, for example, dissociation of hydrogen iodide.

 $2 \text{ HI} \rightarrow \text{H}_2 + \text{I}_2$

Trimolecular or termolecular reactions involve simultaneous collision between three reacting species, for example,

 $2 \text{ NO} + \text{O}_2 \rightarrow 2 \text{NO}_2$

The probability that more than three molecules can collide and react simultaneously is very small. Hence, reactions with the molecularity three are very rare and slow to proceed.

It is, therefore, evident that complex reactions involving more than three molecules in the stoichiometric equation must take place in more than one step.

 $KClO_3 + 6 \ FeSO_4 + 3 \ H_2SO_4 \rightarrow KCl + 3 \ Fe_2(SO_4)_3 + 3 \ H_2O$

This reaction which apparently seems to be of tenth order is actually a second order reaction. This shows that this reaction takes place in several steps. Which step controls the rate of the overall reaction? The question can be answered if we go through the mechanism of reaction, for example, chances to win the relay race competition by a team depend upon the slowest person in the team. Similarly, the overall rate of the reaction is controlled by the slowest step in a reaction called the rate determining step. Consider the decomposition of hydrogen peroxide which is catalysed by iodide ion in an alkaline medium.

 $2H_2O_2 \longrightarrow 2H_2O + O_2$

I-

Alkaline Medium

The rate equation for this reaction is found to be

$$Rate = \frac{-d[H_2O_2]}{dt} = k[H_2O_2] \quad [I^{-}]$$

This reaction is first order with respect to both H_2O_2 and I^- . Evidences suggest that this reaction takes place in two steps:

Step 1: $H_2O_2 + I^- \rightarrow H_2O + IO^-$ Step 2: $H_2O_2 + IO^- \rightarrow H_2O + I^- + O_2$ Both the steps are bimolecular elementary reactions. Species IO⁻ is called as an intermediate since it is formed during the course of the reaction but not in the overall balanced equation. The first step, being slow, is the rate determining step. Thus, the rate of formation of intermediate will determine the rate of this reaction. The detailed description of various steps involved in a reaction is known as the mechanism of a reaction.

We have to keep this in mind that each and every step involved in a reaction proceeds with different rates; only the slowest step of the reaction depicts the overall rate of the reaction. In cases of complex reactions, the molecularity of overall reaction, in general, has no significance.

Thus, from the discussion till now, we conclude the following:

i. Order of the reaction is the sum of the powers of the concentration terms in the rate law expression while molecularity is simply the number of the reacting species.

ii. Order of a reaction is an experimental quantity. It can be zero and even a fraction but molecularity is a theoretical concept and cannot be zero or a non-integer.

iii. Order is applicable to elementary as well as complex reactions whereas molecularity is applicable only for elementary reactions. For complex reaction molecularity has no significance.

iv. For complex reaction, order is given by the slowest step and hence depicts the mechanism of the reaction; on the other hand, molecularity of the slowest step is same as the order of the overall reaction and gives no clue about the mechanism.

v. Order of the reaction depends upon the experimental conditions like temperature and pressure while molecularity of the reaction does not.

Example 4: For a reaction, $A + B \rightarrow$ Product, the rate law is given by, $r = k [A]^{1/2} [B]^2$. What is the order of the reaction?

Solution: The order of the reaction is the sum of powers of the concentration terms in the rate law expression. Here, we have

rate = k $[A]^{1/2} [B]^2$

Therefore, order of the reaction = $\frac{1}{2} + 2 = \frac{5}{2} = 2.5$

Example 5: The conversion of molecules X to Y follows second order kinetics. If concentration of X is increased to three times how will it affect the rate of formation of Y? **Solution:** The reaction $X \rightarrow Y$ follows second order kinetics. Therefore, the rate equation for this reaction will be given as:

Rate = $k [X]^2$

Let the concentration of X, i.e. [X] be 'a' mol L⁻¹, then rate of the reaction is: Rate = $k (a)^2 = k a^2$

If the concentration of X is increased to three times, then $[X] = 3 \times a \mod L^{-1}$

Therefore, rate of the reaction will be:

 $Rate_{new} = k (3a)^2 = 9 k a^2$

Hence, the rate of the reaction or rate of the formation of Y will increase by 9 times.

Example 6: For what type of reactions the rate constant shall have the same units as the rate of the reaction?

Solution: It is for the zero-order reaction.

6. Summary: In first module we have learnt about the rate of the reaction; in this module, we learnt that the number of factors such as temperature, concentration of reactants, catalyst, affect the rate of a reaction. It was observed that with passage of time the concentration of reactant decreases while that of products increases. Also, the rate of the reaction decreases with decrease in concentration of the reactants.

Mathematical representation of rate of a reaction in terms of concentration of the reactants is given by rate law. It has to be determined experimentally and cannot be predicted. The proportionality factor in the rate law is known as rate constant, k. Rate constant and order of a reaction can be determined from rate law or its integrated rate equation.

Order of a reaction with respect to a reactant is the power of its concentration which appears in the rate law equation. The order of a reaction is the sum of all such powers of concentration of terms for different reactants. Order of a reaction can be 0, 1, 2 or any noninteger. Molecularity is defined only for an elementary reaction. Its values are limited from 1 to 3 whereas order can be 0, 1, 2, 3 or even a fraction. Molecularity cannot be zero or a noninteger. Molecularity and order of an elementary reaction are same. Many reactions occur in a number of steps that make the overall mechanism of the reaction and the rate of the reaction is determined by the slowest step of the reaction.