

## 1. Details of Module and its structure

Module Detail	
Subject Name	Chemistry
Course Name	Chemistry 03 (Class XII, Semester 01)
Module Name/Title	Chemical Kinetics : Part 3
Module Id	lech_10403
Pre-requisites	Chemical reaction, concentration terms, rate of reaction, units of rate of reaction, rate law, rate constant, order of the reaction, molecularity of the reaction
Objectives	After learning this module, the student would be able to: <ul style="list-style-type: none"><li>• Derive integrated rate equations for the zero and first order reactions</li><li>• Determine the rate constants for zeroth and first order reactions</li><li>• Determine the half-life of a reaction</li></ul>
Keywords	Rate of the reaction, integrated rate equation, rate constant, order of the reaction, first order reaction, pseudo first order reaction, half- life of a reaction

## 2. Development Team

Role	Name	Affiliation
National MOOC Coordinator (NMC)	Prof. Amarendra P. Behera	CIET, NCERT, New Delhi
Program Coordinator	Dr. Mohd. Mamur Ali	CIET, NCERT, New Delhi
Course Coordinator (CC) / PI	Prof. Alka Mehrotra Prof. R. K. Parashar	DESM, NCERT, New Delhi DESM, NCERT, New Delhi
Course Co-Coordinator / Co-PI	Dr. Aerum Khan	CIET, NCERT, New Delhi
Subject Matter Expert (SME)	Dr. Komal S. Khatri	G. B. Pant Institute of Polytechnic, Okhla Phase II, New Delhi
Review Team	Dr. Amirtha Anand	Dept. of Chemistry, Maitreyi College, New Delhi-110021

---


## TABLE OF CONTENTS

1. Integrated rate equations
2. Zero order reactions
3. First order reactions
4. Half-life of a reaction
5. Pseudo first order reactions
6. Summary

**1. Integrated rate equations:** We have already noted in last module that the concentration dependence of rate is called differential rate equation. It is not always convenient to determine the instantaneous rate, as it is measured by determination of slope of the tangent at point 't' in concentration vs time plot. This makes it difficult to determine the rate law and hence the order of the reaction. In order to avoid this difficulty, we can integrate the differential rate equation to give a relation between directly measured experimental data, i.e., concentrations at different times and rate constant.

The integrated rate equation is used to determine the rate constant of the reaction. Also, it is helpful to calculate the exact time when reaction is either complete or at any intermediate position (i.e. 25% or 50% or 90% completion) and vice versa. The integrated equations are different for the reactions of different reaction orders. We shall determine these equations only for zero and first order chemical reactions.

**2. Zero order reactions:** A reaction with zero order means that the rate of the reaction is proportional to zero power of the concentration of reactants, i.e. independent of the concentration of the reactants. Consider the reaction,



$$\text{Rate} = \frac{-d[R]}{dt} = k[R]^0$$

As any quantity raised to power zero is unity

$$\text{Rate} = \frac{-d[R]}{dt} = k \times 1$$

$$d[R] = -k \cdot dt$$

---

Integrating both sides

$$[R] = -kt + I \quad (1)$$

where I is the constant of integration.

At  $t = 0$ , the concentration of the reactant  $R = [R]_0$ , where  $[R]_0$  is initial concentration of the reactant.

Substituting in equation (1)

$$[R]_0 = -k \times 0 + I$$


$$[R]_0 = I$$

Substituting the value of I in the equation (1)

$$[R] = -kt + [R]_0 \quad (2)$$

Comparing (2) with equation of a straight line,

$y = mx + c$ , if we plot  $[R]$  against  $t$ , we get a straight line (Fig.1) with slope =  $-k$  and intercept equal to  $[R]_0$ .




**Fig. 1.** Variation in the concentration vs time plot for a zero order reaction

(Source: Fig 4.3, page no. 104, Chapter 4: Chemical Kinetics, NCERT Textbook)

Further simplifying equation (2), we get the rate constant,  $k$  as

$$k = \frac{[R]_0 - [R]}{t} \quad (3)$$

All zero order reactions obey the equation (3). If the values for  $[R]_0$ ,  $[R]$  and  $t$  are known then the value of rate constant can be easily evaluated. Zero order reactions are relatively uncommon but they occur under special conditions. Some enzyme catalysed reactions and reactions which occur on metal surfaces are a few examples of zero order reactions. These types of reactions generally occur in heterogeneous systems. For example, the decomposition of gaseous ammonia on a hot platinum surface is a zero order reaction at high pressure.




Rate of the reaction is given as:

$$\text{Rate} = k [\text{NH}_3]^0 = k$$

In this reaction, platinum metal acts as a catalyst. At high pressure, the finely divided metal surface, i.e. platinum surface gets saturated with ammonia gas molecules. So, a further change in reaction conditions is unable to alter the amount of ammonia on the surface of the catalyst making rate of the reaction independent of its concentration. The thermal decomposition of HI on gold surface is another example of zero order reaction.

**3. First order reactions:** In this class of reactions, the rate of the reaction is proportional to the first power of the concentration of the reactant R. For example,



The rate of the reaction is given as:

$$\text{Rate} = \frac{-d[R]}{dt} = k[R]$$

or

$$\frac{-d[R]}{[R]} = kdt$$

Integrating this equation, we get

$$\ln[R] = -kt + I \quad (4)$$

---

Again, I is the constant of integration and its value can be determined easily.

When  $t = 0$ ,  $R = [R]_0$ , where  $[R]_0$  is the initial concentration of the reactant.

Therefore, equation (4) can be written as

$$\ln [R]_0 = -(k \times 0) + I$$

$$\ln [R]_0 = I$$

Substituting the value of I in equation (4), we get

$$\ln [R] = -kt + \ln [R]_0 \quad (5)$$

On rearranging the equation (5), we get

$$\ln \frac{[R]}{[R]_0} = -kt$$

or  $k = \frac{1}{t} \ln \frac{[R]_0}{[R]}$  (6)

At time  $t_1$ , from equation (4)

$$\ln [R]_1 = -kt_1 + \ln [R]_0 \quad (7)$$

At time  $t_2$ ,

$$\ln [R]_2 = -kt_2 + \ln [R]_0 \quad (8)$$

where  $[R]_1$  and  $[R]_2$  are the concentrations of the reactants at time  $t_1$  and  $t_2$  respectively.

Subtracting equation (8) from equation (7),

$$\ln [R]_1 - \ln [R]_2 = -kt_1 - (-kt_2)$$

$$\ln \frac{[R]_1}{[R]_2} = -k(t_1 - t_2) = k(t_2 - t_1)$$

$$k = \frac{1}{(t_2 - t_1)} \ln \frac{[R]_1}{[R]_2} \quad (9)$$


Equation (5) can also be written as

$$\ln \frac{[R]}{[R]_0} = -kt$$

Taking antilog of both sides, we get

$$[R] = [R]_0 e^{-kt} \quad (10)$$

Comparing equation (5) with  $y = mx + c$ , if we plot  $\ln [R]$  against  $t$  (Fig. 2), we get a straight line with slope  $= -k$  and intercept equal to  $\ln [R]_0$ .



**Fig. 2.** A plot between  $\ln [R]$  and  $t$  for a first order reaction.


(Source: Fig. 4, page no. 106, Chapter 4: Chemical Kinetics, NCERT Textbook)

The first order rate equation (6) can also be written in the form

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]} \quad (11)$$

$$\log \frac{[R]_0}{[R]} = \frac{kt}{2.303}$$


If we plot a graph between  $\log [R]_0/[R]$  vs  $t$ , (Fig. 3), the slope  $= (k / 2.303)$ .



**Fig. 3.** A plot of  $\log ([R]_0/[R])$  vs time for a first order reaction.


(Source: Fig 5, page no. 106, Chapter 4: Chemical Kinetics, NCERT Textbook)

Hydrogenation of ethene is an example of first order reaction.



$$\text{Rate} = k [\text{C}_2\text{H}_4]$$


All natural and artificial radioactive decay of unstable nuclei take place by first order kinetics.



$$\text{Rate} = k [\text{Ra}]$$

Decomposition of  $\text{N}_2\text{O}_5$  and  $\text{N}_2\text{O}$  are some more examples of first order reactions.


Let us consider a typical first order gas phase reaction



Let  $p_i$  be the initial pressure of A and  $p_t$  the total pressure at time 't'. Integrated rate equation for such a reaction can be derived as

$$\text{Total pressure, } p_t = p_A + p_B + p_C \text{ (pressure units)}$$

where  $p_A$ ,  $p_B$  and  $p_C$  are the partial pressures of A, B and C, respectively. If  $x$  atm be the decrease in pressure of A at time  $t$  and one mole each of B and C is being formed, the increase in pressure of B and C will also be  $x$  atm each.



At  $t = 0$ ,  $p_i$  atm      0 atm      0 atm

At time  $t$ ,  $(p_i - x)$  atm       $x$  atm       $x$  atm

where  $p_i$  is the initial pressure at time  $t = 0$ .

$$p_t = (p_i - x) + x + x = p_i + x$$

$$x = (p_t - p_i)$$


where  $p_A = p_i - x = p_i - (p_t - p_i)$

$$= 2 p_i - p_t$$

$$k = \left( \frac{2.303}{t} \right) \left( \log \frac{p_i}{p_A} \right) \quad (12)$$

$$\left( \frac{2.303}{t} \right) \log \frac{p_i}{(2 p_i - p_t)}$$

**Example 1:** The initial concentration of  $N_2O_5$  in the following first order reaction



was  $1.24 \times 10^{-2} \text{ mol L}^{-1}$  at 318 K. The concentration of  $N_2O_5$  after 60 minutes was  $0.20 \times 10^{-2} \text{ mol L}^{-1}$ . Calculate the rate constant of the reaction at 318 K.

**Solution:** For the first order reaction, we have

$$\log \frac{[R]_1}{[R]_2} = \frac{k(t_2 - t_1)}{2.303}$$

$$k = \frac{2.303}{(t_2 - t_1)} \log \frac{[R]_1}{[R]_2}$$




$$\frac{2.303}{(60 \text{ min} - 0 \text{ min})} \log \frac{1.24 \times 10^{-2} \text{ mol L}^{-1}}{0.20 \times 10^{-2} \text{ mol L}^{-1}}$$

$$\frac{2.303}{60} \log 6.2 \text{ min}^{-1}$$

$$k = 0.0304 \text{ min}^{-1}$$


**Example 2:** The following data were obtained during the first order thermal decomposition of  $\text{N}_2\text{O}_5$  (g) at constant volume:



S.No.	Time/s	Total Pressure/(atm)
1.	0	0.5
2.	100	0.512

Calculate the rate constant.

**Solution:** Let the pressure of  $\text{N}_2\text{O}_5$  (g) decrease by  $2x$  atm. As two moles of  $\text{N}_2\text{O}_5$  decompose to give two moles of  $\text{N}_2\text{O}_4$  (g) and one mole of  $\text{O}_2$  (g), the pressure of  $\text{N}_2\text{O}_4$  (g) increases by  $2x$  atm and that of  $\text{O}_2$  (g) increases by  $x$  atm.



At  $t = 0$ ,            0.5 atm                    0 atm                    0 atm

At time  $t$ ,         $(0.5 - 2x)$  atm             $2x$  atm                     $x$  atm

$$p_t = p_{\text{N}_2\text{O}_5} + p_{\text{N}_2\text{O}_4} + p_{\text{O}_2}$$

$$= (0.5 - 2x) + 2x + x = 0.5 + x$$

$$= x = p_t - 0.5$$

$$p_{\text{N}_2\text{O}_5} = 0.5 - 2x = 0.5 - 2(p_t - 0.5) = 1.5 - 2p_t$$

At  $t = 100$  s,  $p_t = 0.512$  atm

---

$$p_{N_2O_5} = 1.5 - 2 \times 0.512 = 0.476 \text{ atm}$$

Using equation (12)

$$k = \left( \frac{2.303}{t} \right) \left( \log \frac{p_i}{p_A} \right)$$

$$k = \left( \frac{2.303}{100 \text{ s}} \right) \left( \log \frac{0.5 \text{ atm}}{0.476 \text{ atm}} \right) = \frac{2.303}{100 \text{ s}} \times 0.0216 = 4.98 \times 10^{-4} \text{ s}^{-1}$$

**Example 3:** For a first order reaction, calculate the time required to reduce the concentration of a reactant from 10 g to 3 g. (Given: the first order reaction rate constant is  $2.51 \times 10^{-3} \text{ s}^{-1}$ )

**Solution:** Initial concentration of reactant,  $[R]_0 = 10 \text{ g}$

Final concentration of reactant,  $[R]_t = 3 \text{ g}$

Rate constant =  $2.51 \times 10^{-3} \text{ s}^{-1}$

For a first order reaction, we have

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$

or

$$\frac{10}{3} = \frac{2.303}{2.51 \times 10^{-3}} \times 0.523 \text{ s}$$
$$t = \frac{2.303}{k} \log \frac{[R]_0}{[R]} = \frac{2.303}{2.51 \times 10^{-3}} \log$$

$$t = 0.48 \times 10^3 \text{ s}$$

**4. Half-life of a reaction:** The half-life of a reaction is the time in which the concentration of a reactant is reduced to one half of its initial concentration. In other words, half-life is the time required for the completion of half of the reaction. It is represented as  $t_{1/2}$ .

Half-Life for zero order reaction: For a zero order reaction, rate constant is given as:

$$k = \frac{[R]_0 - [R]}{t}$$

At  $t = t_{1/2}$ ,  $[R] = [R]_0/2$

Hence, the rate constant at  $t_{1/2}$  becomes

$$k = \frac{[R]_0 - \left(\frac{1}{2}[R]_0\right)}{t_{1/2}}$$

$$t_{1/2} = \frac{[R]_0}{2k}$$

It is clear that  $t_{1/2}$  for a zero order reaction is directly proportional to the initial concentration of the reactants and inversely proportional to the rate constant.

Half-Life for first order reaction: For the first order reaction we,

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]} \quad (13)$$

At  $t = t_{1/2}$ ,  $[R] = [R]_0/2$

S.No.	Time/s	Total Pressure/(atm)
1.	0	0.5
2.	100	0.512

Hence, the rate constant at  $t_{1/2}$  becomes

$$k = \frac{2.303}{t_{1/2}} \log \frac{[R]_0}{\left(\frac{1}{2}[R]_0\right)} \quad (14)$$

$$t_{1/2} = \frac{2.303}{k} \log 2 = \left(\frac{2.303}{k}\right) \times 0.301$$

$$t_{1/2} = \frac{0.693}{k} \quad (15)$$

---

It can be seen that for a first order reaction, half-life period is constant, i.e., it is independent of initial concentration of the reacting species. The half-life of a first order equation is readily calculated from the rate constant and vice versa.

**For zero order reaction  $t_{1/2} \propto [R]_0$ . For first order reaction  $t_{1/2}$  is independent of  $[R]_0$ .**

**Example 4:** A first order reaction is found to have a rate constant,  $k = 5.5 \times 10^{-14} \text{ s}^{-1}$ . Find the half-life of the reaction.

**Solution:** Half-life for a first order reaction is

$$t_{1/2} = \frac{0.693}{k}$$

$$t_{1/2} = \frac{0.693}{5.5 \times 10^{-14} \text{ s}^{-1}}$$

$$t_{1/2} = 1.26 \times 10^{13} \text{ s}$$

**Example 5:** Show that in a first order reaction, time required for completion of 99.9% is 10 times of half-life ( $t_{1/2}$ ) of the reaction.

**Solution:** When reaction is completed 99.9%,  $[R] = [R]_0 - 0.999[R]_0$

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]_0 - 0.999[R]_0}$$

$$k = \frac{2.303}{t} \log 10^3$$

$$t = \frac{6.909}{k}$$

For half-life of the reaction, we know

$$t_{1/2} = \frac{0.693}{k}$$

$$\frac{t}{t_{1/2}} = \frac{6.909}{k} \times \frac{k}{0.693} = 10$$

---

**Example 6:** The half-life period of a first order reaction is 200 sec. Calculate the rate constant for this reaction.

**Solution:** For a first order reaction, half-life is given as

$$t_{1/2} = \frac{0.693}{k}$$

Therefore,

$$k = \frac{0.693}{t_{1/2}}$$

We have,  $t_{1/2} = 200$  sec

$$k = \frac{0.693}{200\text{s}} = 3.465 \times 10^{-3} \text{ s}^{-1}$$

**Example 7:** The half-life period for a first order reaction is 45 minutes. Calculate the percentage of the reactant left after 180 minutes.

**Solution:** For a first order reaction, it is given that

$$t_{1/2} = 45 \text{ min}$$

The rate constant for the reaction will be

$$k = \frac{0.693}{t_{1/2}} = \frac{0.693}{45 \text{ min}} = 1.54 \times 10^{-2} \text{ min}^{-1}$$

The rate expression for first order is given by

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]} \quad \text{or} \quad \frac{[R]_0}{[R]} = \exp(kt) \quad \text{or} \quad \frac{[R]}{[R]_0} = \exp(-kt)$$

At  $t = 180$  min

$$\frac{[R]}{[R]_0} = \exp(-1.54 \times 10^{-2} \times 180) = 0.0625$$

Therefore, after 180 minutes only 6.25% of the reactant is left.

**Example 8:** For a zero order reaction, the rate constant is  $0.14 \text{ mol L}^{-1} \text{ s}^{-1}$ . If after two minutes the concentration of the reactant is  $0.15 \text{ mol L}^{-1}$ , calculate its initial concentration.

**Solution:** For a zero order reaction we have,

$$k = \frac{[R]_0 - [R]}{t}$$

Given:  $k = 0.14 \text{ mol L}^{-1} \text{ s}^{-1}$ ;  $[R] = 0.15 \text{ mol L}^{-1}$ ;  $t = 2 \text{ min} = 2 \times 60 \text{ sec} = 120 \text{ sec}$

$$[R]_0 = (k \times t) + [R]$$

$$= \{(0.14 \times 120) + 0.15\} \text{ mol L}^{-1}$$

$$= (16.8 + 0.15) \text{ mol L}^{-1}$$

$$= 16.95 \text{ mol L}^{-1}$$

Table 1 summarises the mathematical features of integrated laws of zero and first order reactions.

**Table 1.** Integrated Rate Laws for the Reactions of Zero and First Order Reactions

Order	Reaction type	Differential rate law	Integrated rate law	Straight line plot	Half-Life	Units of $k$
0	$R \rightarrow P$	$\frac{d[R]}{dt} = -k$	$kt = [R]_0 - [R]$	$[R]$ vs $t$	$[R]_0 / 2k$	conc $\text{time}^{-1}$ or $\text{mol L}^{-1} \text{ s}^{-1}$
1	$R \rightarrow P$	$\frac{d[R]}{dt} = -k[R]$	$[R] = [R]_0 \exp(-kt)$ or $kt = \ln\{[R]_0/[R]\}$	$\ln[R]$ vs $t$	$\ln 2 / k$ or $0.693 / k$	$\text{time}^{-1}$ or $\text{s}^{-1}$

**5. Pseudo first order reactions:** The order of a reaction is sometimes altered by conditions. Consider a chemical reaction between two substances when one reactant is present in large excess. During the hydrolysis of 0.01 mol of ethyl acetate with 10 mol of water, amounts of the various constituents at the beginning ( $t = 0$ ) and at the completion ( $t$ ) of the reaction are given as under.





---

$$\text{Rate} = k [\text{CH}_3\text{COOCH}_3][\text{H}_2\text{O}]$$

**Solution:** For pseudo first order reaction, the reaction should be first order with respect to ester when  $[\text{H}_2\text{O}]$  is constant. The rate constant  $k'$  for pseudo first order reaction is

$$k' = \frac{2.303}{t} \log \frac{C_0}{C}$$

where  $k' = k [\text{H}_2\text{O}]$

From the above data we note

$t / \text{min}$	$C / \text{mol L}^{-1}$	$k' / \text{min}^{-1}$
0	0.8500	-
30	0.8004	$2.004 \times 10^{-3}$
60	0.7538	$2.002 \times 10^{-3}$
90	0.7096	$2.005 \times 10^{-3}$

It can be seen that  $k' = k[\text{H}_2\text{O}]$  is constant and equal to  $2.004 \times 10^{-3} \text{ min}^{-1}$  and hence, it is pseudo first order reaction. We can now determine  $k$  from

$$k [\text{H}_2\text{O}] = 2.004 \times 10^{-3} \text{ min}^{-1}$$

$$k [55 \text{ mol L}^{-1}] = 2.004 \times 10^{-3} \text{ min}^{-1}$$

$$k = 3.64 \times 10^{-5} \text{ mol}^{-1} \text{ L min}^{-1}$$

**Example 10:** In a pseudo first order hydrolysis of ester in water the following results were obtained:

$t / \text{min}$	0	30	60	90
$[\text{Ester}] / \text{M}$	0.55	0.31	0.17	0.085

- Calculate the average rate of the reaction between the time interval 30 to 60 seconds.
- Calculate the pseudo first order rate constant for the hydrolysis of ester.



**Solution:** a) Average rate of the reaction between time interval 30 to 60 seconds is given by:

$$\text{Average rate} = \frac{C_2 - C_1}{\Delta t} = \frac{(0.17 - 0.31)}{(60 - 30)} = \frac{-0.14}{30} = -4.67 \times 10^{-3} \text{ M s}^{-1}$$

Negative sign indicates that as the concentration of ester decreases with passage of time, the rate of reaction also decreases with time.

b) Pseudo first order rate constant is given as

$$k = \frac{2.303}{t} \log \frac{a}{a-x}$$

where a: initial concentration and (a - x): concentration after time t.

Here, a = 0.55 M. The rate constant at different instant of time is tabulated below:

t/ min	C/ mol L <sup>-1</sup>	k'/ s <sup>-1</sup>
30	0.31	1.91 × 10 <sup>-2</sup>
60	0.17	1.95 × 10 <sup>-2</sup>
90	0.085	2.06 × 10 <sup>-2</sup>

The values of rate constant at every instant of time is almost the same which confirms that the reaction is of first order. The actual value for rate constant will be:

$$k = \frac{(1.91 + 1.95 + 2.06)}{3} \times 10^{-2} \text{ s}^{-1} = 1.97 \times 10^{-2} \text{ s}^{-1}$$

**6. Summary:** In this module we have learnt about the integrated forms of the rate law equations for zero and first order reactions. A reaction with zero order means that the rate of the reaction is proportional to zero power of the concentration of reactants. In first order reactions (like R → P), the rate of the reaction is proportional to the first power of the concentration of the reactant R. The integrated rate equations for zero and first order reactions were derived. The half-life of a reaction is the time in which the concentration of a reactant is reduced to one half of its initial concentration. For zero and first order reactions, the half-life periods were derived. There are also some reactions called pseudo first order reactions. These reactions appear to be of higher order but actually follow first order kinetics.